

HADRONIC DECAYS OF BARYONS IN POINT-FORM RELATIVISTIC QUANTUM MECHANICS

T. MELDE*, W. PLESSAS, AND R. F. WAGENBRUNN

*Theoretische Physik, Institut für Physik, Universität Graz
Universitätsplatz 5, A-8010 Graz, Austria*

L. CANTON

*INFN, Sezione di Padova, and Dipartimento di Fisica dell'Università di Padova
Via F. Marzolo 8, I-35131 Padova, Italy*

We discuss strong decays of baryon resonances within the concept of relativistic constituent quark models. In particular, we follow a Poincaré-invariant approach along the point form of relativistic quantum mechanics. Here, we focus on pionic decay modes of N and Δ resonances. It is found that the covariant quark-model predictions calculated in the point-form spectator model in general underestimate the experimental data considerably. This points to a systematic defect in the used decay operator and/or the baryon wave functions. From a detailed investigation of the point-form decay operator it is seen that the requirement of translational invariance implies effective many-body contributions. Furthermore, one has to employ a normalization factor in the definition of the decay operator in the point-form spectator model. Our analysis suggests that this normalization factor is best chosen consistently with the one used for the electromagnetic and axial current operators for elastic nucleon form factors.

1. Introduction

Constituent quark models (CQMs) provide an effective description of hadrons in the low-energy regime of quantum chromodynamics (QCD). It has been of particular interest to find an appropriate type of (phenomenological) interaction between the constituent quarks, which is usually comprised of a confinement and a hyperfine part. Usually the hyperfine interaction is derived from one-gluon exchange (OGE)¹. Beyond that also alternative types of CQMs have been suggested such as the ones based on instanton-induced (II) forces² or Goldstone-boson-exchange

*Work partially supported by INFN and MIUR-PRIN.

(GBE) dynamics³. All these variants of modern CQMs describe the overall trends of the hadronic spectra reasonably well. With regard to baryons, however, only the GBE CQM succeeds in reproducing simultaneously the level ordering of positive- and negative-parity resonances as well as the $N - \Delta$ splitting in agreement with phenomenology⁴. The typical spectra of the different types of CQMs are exemplified in fig. 1.

Another problem has been the role of relativity. Specifically, it has been found that covariant predictions of relativistic CQMs for electroweak nucleon form factors agree surprisingly well with experimental data. This is especially true for the II CQM in a Bethe-Salpeter type approach⁷. The same has been found for the GBE CQM and likewise the OGE CQM within the point-form approach of relativistic quantum mechanics^{8,9,10}. In contrast, any nonrelativistic calculation fails drastically.

The covariant results obtained so far for mesonic decay widths of N and Δ resonances have shown a systematic underestimation of the experimental data for any of the three types of CQMs^{11,12,13}. Clearly, this points to shortcomings in the baryon (resonance) wave functions and/or the decay operators employed. In this contribution we give a review of the current status of the investigations on hadronic decays of N and Δ resonances.




Figure 1. Nucleon excitation spectra of three different types of relativistic CQMs. The left panel shows the nucleon spectrum and the right panel the Δ spectrum. In each column the left horizontal lines represent the results of the relativized Bhaduri-Cohler-Nogami CQM⁵, the middle ones of the II CQM (Version A)², and the right ones of the GBE CQM³. The shadowed boxes give the experimental data with their uncertainties after the latest compilation of the PDG⁶.

2. Theory

Generally, the decay width Γ of a resonance is defined by the expression

$$\Gamma = 2\pi\rho_f |F(i \rightarrow f)|^2, \quad (1)$$

where $F(i \rightarrow f)$ is the transition amplitude and ρ_f is the phase-space factor. In eq. (1) one has to average over the initial and to sum over the final spin-isospin projections. A common problem in nonrelativistic approximations of the transition amplitude is the ambiguity of the proper phase-space factor^{14,15,16}. Here, we present a Poincaré-invariant approach, adhering to the point form of relativistic quantum mechanics¹⁷. In this case the generators of the Lorentz transformations remain purely kinematical and the theory is manifestly covariant¹⁸. This also allows to resolve the ambiguity in the phase-space factor. The interactions between the constituent quarks are introduced into the (invariant) mass operator following the Bakamjian-Thomas construction¹⁹. In this approach the covariant transition amplitude for mesonic decays is defined via the matrix element of the decay operator

$$\begin{aligned} F(i \rightarrow f) &= \langle P', J', \Sigma' | \hat{D}_m | P, J, \Sigma \rangle \\ &= \frac{2}{MM'} \sum_{\sigma_i \sigma'_i} \sum_{\mu_i \mu'_i} \int d^3\vec{k}_2 d^3\vec{k}_3 d^3\vec{k}'_2 d^3\vec{k}'_3 \\ &\quad \sqrt{\frac{(\omega_1 + \omega_2 + \omega_3)^3}{2\omega_1 2\omega_2 2\omega_3}} \sqrt{\frac{(\omega'_1 + \omega'_2 + \omega'_3)^3}{2\omega'_1 2\omega'_2 2\omega'_3}} \\ &\quad \Psi_{M'J'\Sigma'}^* \left(\vec{k}'_i; \mu'_i \right) \prod_{\sigma'_i} D_{\sigma'_i \mu'_i}^{\star \frac{1}{2}} \{ R_W [k'_i; B(V')] \} \\ &\quad \langle p'_1, p'_2, p'_3; \sigma'_1, \sigma'_2, \sigma'_3 | \hat{D}_m | p_1, p_2, p_3; \sigma_1, \sigma_2, \sigma_3 \rangle \\ &\quad \prod_{\sigma_i} D_{\sigma_i \mu_i}^{\frac{1}{2}} \{ R_W [k_i; B(V)] \} \Psi_{MJ\Sigma} \left(\vec{k}_i; \mu_i \right), \end{aligned} \quad (2)$$

where overall momentum conservation, $P_\mu - P'_\mu = Q_\mu$, is explicitly satisfied; Q_μ being the meson four-momentum. Here $|P, J, \Sigma\rangle$ and $\langle P', J', \Sigma'|$ are the eigenstates of the decaying resonance and the outgoing nucleon ground state, respectively. The eigenstates are denoted by the eigenvalues P, J, Σ of the four-momentum operator $\hat{P}$, the total angular-momentum operator $\hat{J}$ and its z -component $\hat{\Sigma}$. The corresponding rest-frame baryon wave functions $\Psi_{M'J'\Sigma'}^*$ and $\Psi_{MJ\Sigma}$ stem from the velocity-state representations of the baryon states $\langle P', J', \Sigma'|$ and $|P, J, \Sigma\rangle$. The rest-frame quark momenta $\vec{k}_i$, for which $\sum_i \vec{k}_i = \vec{0}$, are related to the individual quark four-momenta by

the Lorentz boost relations $p_i = B(v) k_i$, with analogous relations holding for the primed variables.

In previous studies of the electroweak nucleon structure^{8,9,10} a point-form spectator model (PFSM) for the electromagnetic and axial currents performed very well. Consequently, in a first investigation of mesonic decays we adopt a PFSM also for the decay operator. Assuming a pseudovector quark-meson coupling we express it by

$$\begin{aligned} \langle p'_1, p'_2, p'_3; \sigma'_1, \sigma'_2, \sigma'_3 | \hat{D}_m | p_1, p_2, p_3; \sigma_1, \sigma_2, \sigma_3 \rangle = \\ 3N \frac{ig_{qq\pi}}{2m_1 (2\pi)^{\frac{3}{2}}} \bar{u}(p'_1, \sigma'_1) \gamma_5 \gamma^\mu \lambda_m u(p_1, \sigma_1) Q_\mu \\ 2p_{20} \delta(\vec{p}_2 - \vec{p}'_2) 2p_{30} \delta(\vec{p}_3 - \vec{p}'_3) \delta_{\sigma_2 \sigma'_2} \delta_{\sigma_3 \sigma'_3}, \end{aligned} \quad (3)$$

with the flavor matrix λ_m characterizing the particular decay mode. Here, the incoming and outgoing momenta of the active quark are determined uniquely by the overall momentum conservation of the transition amplitude, $P_\mu - P'_\mu = Q_\mu$, together with the two spectator conditions. Generally, the momentum transferred to the active quark, $\vec{p}_1 - \vec{p}'_1 = \vec{q}$, is different from the momentum transfer $\vec{Q}$ to the baryon as a whole. It has been shown that this is a consequence of translational invariance and also induces effective many-body contributions in the definition of the spectator-model decay operator²⁰. Furthermore, in eq. (3) there appears an overall normalization factor N . In the PFSM electromagnetic and axial currents for the nucleon elastic form factors the factor

$$N = \left(\frac{M}{\sum_i \omega_i} \frac{M'}{\sum_i \omega'_i} \right)^{\frac{3}{2}} \quad (4)$$

was employed to recover the proper proton charge^{8,9,10}. It depends on the individual quark momenta through the ω_i and the on-mass-shell conditions of the quarks. However, this is not a unique choice, if only Poincaré invariance and charge normalization are imposed. Also any other normalization factor of the asymmetric form

$$N(y) = \left(\frac{M}{\sum_i \omega_i} \right)^{3y} \left(\frac{M'}{\sum_i \omega'_i} \right)^{3(1-y)} \quad (5)$$

would be possible²¹. Below we also discuss the consequences of these further choices of N with regard to the pionic decay widths.

3. Results

In table 1 we quote the covariant predictions of the GBE and OGE CQMs for pionic decay widths calculated with the PFSM decay operator of eq. (3) with the normalization factor of eq. (4). For comparison, we also included the results obtained with the II CQM along the Bethe-Salpeter approach²². It is apparent that the results show a systematic underestimation of the experimental data. Only the N_{1535}^* and N_{1710}^* predictions agree with the experimental values.

We also present the numerical results for the decay widths with specific attention to the $\Delta\pi$ branching ratios, for the magnitudes of which a striking relationship is found to the underestimation of the experimental value: the larger the $\Delta\pi$ branching ratio of a resonance, the smaller the relative prediction for the π decay width. At this point it is not yet clear, if this behaviour is purely accidental or substantiates a shortcoming of the present description of baryon decays. In particular, additional degrees of freedom might be missing.

Let us now examine the influence of the normalization factor N in eq. (3). In fig. 2 we show the results for different possible choices of $N(y)$

Table 1. PFSM predictions for π decay widths of the relativistic GBE³ and OGE⁵ CQMs in comparison to the Bethe-Salpeter results²² of the II CQM² and experimental data⁶. In the last three columns the theoretical results are expressed as percentage fractions of the (best-estimate) experimental values in order to be compared to the measured $\Delta\pi$ branching ratios.

Decays	Experiment	Rel. CQM			$\Delta\pi$	% of Exp. Width		
$\rightarrow N\pi$	[MeV]	GBE	OGE	II	branching ratio	GBE	OGE	II
N_{1440}^*	$(227 \pm 18)_{-59}^{+70}$	33	53	38	20 – 30%	14	24	17
N_{1520}^*	$(66 \pm 6)_{-5}^{+9}$	17	16	38	15 – 25%	26	24	58
N_{1535}^*	$(67 \pm 15)_{-17}^{+28}$	90	119	33	< 1%	134	178	49
N_{1650}^*	$(109 \pm 26)_{-3}^{+36}$	29	41	3	1 – 7%	27	38	3
N_{1675}^*	$(68 \pm 8)_{-4}^{+14}$	5.4	6.6	4	50 – 60%	8	10	6
N_{1700}^*	$(10 \pm 5)_{-3}^{+3}$	0.8	1.2	0.1	> 50%	8	12	1
N_{1710}^*	$(15 \pm 5)_{-5}^{+30}$	5.5	7.7	<i>n/a</i>	15 – 40%	37	51	<i>n/a</i>
Δ_{1232}	$(119 \pm 1)_{-5}^{+5}$	37	32	62	–	31	27	52
Δ_{1600}	$(61 \pm 26)_{-10}^{+26}$	0.07	1.8	<i>n/a</i>	40 – 70%	≈ 0	3	<i>n/a</i>
Δ_{1620}	$(38 \pm 8)_{-6}^{+8}$	11	15	4	30 – 60%	29	39	11
Δ_{1700}	$(45 \pm 15)_{-10}^{+20}$	2.3	2.3	2	30 – 60%	5	5	4




Figure 2. Dependence of the π decay widths on the asymmetry parameter y in the normalization factor of eq. (5) for selected N and Δ resonances.

according to eq. (5) in case of the GBE CQM. It is seen that in the range $0 \leq y \leq 1$ all decay widths grow rapidly with increasing y . Thus one could enhance or reduce the results as compared to the ones obtained with the symmetric factor $N(y = \frac{1}{2})$. While all other choices of $N(y)$ are also allowed, the symmetric factor was used for the electroweak nucleon form factors in refs.^{8,9,10} and likewise the results given in table 1. We consider this as the best choice also for the baryon decays. It has the property that no theoretical result overshoots the corresponding experimental value. This is welcome and can be considered as reasonable, since one may expect further contributions not yet included in the present decay operator to raise the theoretical predictions.

The normalization factor N in eq. (3) effectively introduces a momentum cut-off. This can even better be seen if we investigate the form

$$N(x) = \left(\frac{M}{\sum_i \omega_i} \frac{M'}{\sum_i \omega'_i} \right)^{\frac{x}{2}} \quad (6)$$

with an arbitrary exponent x . It still represents a Poincaré-invariant construction but it does not guarantee for the proper charge normalization unless $x = 3$. It is clearly seen in fig. 3 that the choice $x = 0$ representing the bare case without normalization factor yields unreasonable results. The theoretical decay widths then evolve smoothly with increasing exponent x . For certain resonances (namely, N_{1440}^* , N_{1710}^* , and Δ_{1600}) the decay widths have a minimum. We notice that these are known as so-called structure-dependent resonances²³. They are the radial excitations of the N and Δ ground states, respectively, with a corresponding nodal behaviour in their wave functions. These characteristics are quite distinct from the other reso-




Figure 3. Dependence of the π decay widths on the exponent x of the normalization factor in eq. (6) for selected N and Δ resonances.

nances, which show a monotonous dependence on the exponent x . Through this study we get interesting insights into the causes for occurrence of narrow decay widths, at least for baryon states with nodal behaviour.

If we assume again the criterion that the theoretical predictions for decay widths with the decay operator (3) should not exceed the experimental data, we find that the case with $x = 3$ (and $y = \frac{1}{2}$) is the optimal choice. This leads to a single postulate for normalizing the PFSM operators for strong and electroweak processes.

4. Summary

We have discussed a Poincaré-invariant description of strong baryon resonance decays in point-form relativistic quantum mechanics. Covariant predictions of relativistic CQMs have been shown for π decay widths. They are considerably different from previous nonrelativistic results or results with relativistic corrections included. The covariant results calculated with a spectator-model decay operator show a uniform trend. In almost all cases the corresponding theoretical predictions underestimate the experimental data considerably. This is true in the framework of Poincaré-invariant quantum mechanics (here in point form) as well as in the Bethe-Salpeter approach²². Indications have been given that for a particular resonance the size of the underestimation is related to the magnitude of the $\Delta\pi$ branching ratio. This hints to systematic shortcomings in the description of the decay widths.

The investigation of different possible choices for the normalization factor occurring in the spectator-model decay operator has led to the sugges-

tion that the symmetric choice is the most natural one. It is also consistent with the same (symmetric) choice adopted before for the spectator-model currents in the study of the electroweak nucleon form factors.

Acknowledgements

This work was supported by the Austrian Science Fund (Project P16945). T. M. would like to thank the INFN and the Physics Department of the University of Padova for their hospitality and MIUR-PRIN for financial support.

References

1. S. Capstick and N. Isgur, Phys. Rev. D **34**, 2809 (1986).
2. U. Loering, B. C. Metsch, and H. R. Petry, Eur. Phys. J. **A10**, 395 (2001); *ibid.* 447 (2001).
3. L. Y. Glozman, W. Plessas, K. Varga, and R. F. Wagenbrunn, Phys. Rev. D **58**, 094030 (1998).
4. L. Y. Glozman *et al.*, Phys. Rev. C **57**, 3406 (1998).
5. L. Theussl, R. F. Wagenbrunn, B. Desplanques, and W. Plessas, Eur. Phys. J. **A12**, 91 (2001).
6. S. Eidelman *et al.*, Phys. Lett. **B592**, 1 (2004).
7. D. Merten *et al.*, Eur. Phys. J. **A14**, 477 (2002).
8. R. F. Wagenbrunn *et al.*, Phys. Lett. **B511**, 33 (2001).
9. L. Y. Glozman *et al.*, Phys. Lett. **B516**, 183 (2001).
10. S. Boffi *et al.*, Eur. Phys. J. **A14**, 17 (2002).
11. T. Melde, W. Plessas, and R. F. Wagenbrunn, Few-Body Syst. Suppl. **14**, 37 (2003).
12. B. Metsch, U. Loring, D. Merten, and H. Petry, Eur. Phys. J. **A18**, 189 (2003).
13. T. Melde, W. Plessas, and R. F. Wagenbrunn, Contribution to the N*2004 Workshop, Grenoble, to appear in the Proceedings; hep-ph/0406023 (2004).
14. P. Geiger and E. S. Swanson, Phys. Rev. D **50**, 6855 (1994).
15. S. Kumano and V. R. Pandharipande, Phys. Rev. D **38**, 146 (1988).
16. R. Kokoski and N. Isgur, Phys. Rev. D **35**, 907 (1987).
17. B. D. Keister and W. N. Polyzou, Adv. Nucl. Phys. **20**, 225 (1991).
18. W. H. Klink, Phys. Rev. C **58**, 3587 (1998).
19. B. Bakamjian and L. H. Thomas, Phys. Rev. **92**, 1300 (1953).
20. T. Melde, L. Canton, W. Plessas, and R. F. Wagenbrunn, hep-ph/0411322 (2004).
21. T. Melde, L. Canton, W. Plessas, and R. F. Wagenbrunn, Contribution to the Mini-Workshop on Quark Dynamics, Bled, to appear in the Proceedings; hep-ph/0410274 (2004).
22. B. Metsch, Contribution to the 10th International Conference on Hadron Spectroscopy, Aschaffenburg, 2003; hep-ph/0403118 (2004).
23. R. Koniuk and N. Isgur, Phys. Rev. D **21**, 1868 (1980).